

Guideline



CCHMC Trauma Service Operation Guidelines

Title: Venous Thromboembolism (VTE) Prophylaxis

Effective Date: 06/2018

Number: TR-23

Page: 1 of 5

1.0 SCOPE

- 1.1 Care of the Trauma Services Patient at CCHMC.

2.0 DEFINITIONS

- 2.1. **Altered mobility:** A permanent or temporary state in which the child has a limitation in independent, purposeful physical movement of the body or of one or more extremities.
- 2.2. **Deep Vein Thrombosis (DVT):** A blood clot (thrombus) that was initially formed in a deep (non-peripheral) vein.
- 2.3. **Graduated Compression Stocking (GCS):** Elastic stockings, either knee- or thigh-high, also known as TED hose.
- 2.4. **Risk category: Refer to VTE Risk Factors algorithm**
 - 2.4.1. **Low risk:** No VTE risk factors
 - 2.4.2. **Moderate risk:** Multiple risk factors for VTE in the absence of altered mobility or has altered mobility with one or fewer additional risk factors.
 - 2.4.3. **High risk:** Altered mobility plus two or more additional risk factors
- 2.5. **Sequential Compression Device (SCD):** A device designed to intermittently squeeze blood from underlying deep veins in the leg upon compression of an inflatable sleeve, and to allow the blood to flow again when it decompresses.
- 2.6. **Venous Thromboembolism (VTE):** A blood clot (thrombus) in a vein or one that has broken free and is carried in the bloodstream (embolus).

3.0 GUIDELINE

- 3.1. It is recommended that patients age 10 – 17 years be assessed for VTE risk factors, and based on that assessment, assigned to a risk category (low, moderate, high).
 - 3.1.1. At the time of inpatient admission; and
 - 3.1.2. Reassessed at 48 – 72 hours of hospitalization.
 - 3.1.3. This should be documented in the patient's medical record.
- 3.2. It is recommended that VTE prophylaxis be administered based on risk category as soon as feasible, but within 24 hours of assessment, unless there are contraindications (See algorithm).
- 3.3. If planning to initiate pharmacologic prophylaxis it is recommended:
 - 3.3.1. In surgical patients to seek surgical input regarding bleeding risk prior to initiation
 - 3.3.2. See [BES# 049](#) for management of LMWH prophylaxis
 - 3.3.3. Obtain Hematology consultation when considering alternative pharmacologic agents.
- 3.4. Refer to [BES# 181](#) algorithm below for Risk Category Assessment and Prophylaxis for VTE.

Risk Assessment for Thrombosis for Non-Bariatric Surgery Patients ≥ 10 years old

RN or MD/APN to Complete:

<p>Risk factors?</p>	<p>History</p> <ul style="list-style-type: none"> <input type="checkbox"/> Personal history of blood clot <input type="checkbox"/> Family history of clotting disorder or clots <p>High-risk Medical conditions</p> <ul style="list-style-type: none"> <input type="checkbox"/> Known clotting disorder <input type="checkbox"/> Blood stream infection (currently on antibiotics for positive blood culture) <input type="checkbox"/> Cancer <input type="checkbox"/> Chronic inflammatory condition (i.e. Crohn’s, Ulcerative Colitis, Lupus) <input type="checkbox"/> Nephrotic syndrome <input type="checkbox"/> Trauma patient if > 1 lower extremity (LE) fracture, pelvic fracture, or spinal cord injury <p>Medications</p> <ul style="list-style-type: none"> <input type="checkbox"/> Estrogen (i.e. birth control) in past 2 months (Depo shots don’t contain estrogen) <p>Physical Exam</p> <ul style="list-style-type: none"> <input type="checkbox"/> Obesity (BMI > 95th percentile, see growth chart in Epic) <input type="checkbox"/> PICC or central line 	<p>Total number of risk factors:</p> <p>_____</p>
<p>Contraindications to SCD (sequential compression device)?</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Current DVT <input type="checkbox"/> Fracture of lower extremity (LE) <input type="checkbox"/> Skin conditions affecting LE (burn, dermatitis, wound, epidermolysis bullosa) 	<ul style="list-style-type: none"> <input type="checkbox"/> Yes (if any checked) <input type="checkbox"/> No
<p>Contraindications to Lovenox?</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Active bleeding <input type="checkbox"/> Known bleeding disorder <input type="checkbox"/> Epidural or lumbar puncture in the last 12 hours <input type="checkbox"/> Platelets <50,000/mm or heparin-induced thrombocytopenia <input type="checkbox"/> Brain tumor <input type="checkbox"/> Pelvic fracture in last 48 hours <input type="checkbox"/> Recent or scheduled neurosurgical procedure within 48 hours <input type="checkbox"/> Uncontrolled hypertension 	<ul style="list-style-type: none"> <input type="checkbox"/> Yes (if any checked) <input type="checkbox"/> No

- See next page for page 2 of risk assessment

For MD/APN to complete:

<p>≥2 Risk Factors</p>	<p>Anticipated altered mobility > 48 hours?</p> <ul style="list-style-type: none"> <input type="checkbox"/> Bedrest or significant activity restriction <input type="checkbox"/> Any line or tube that restricts mobility (i.e. epidural, foley, NG to continuous suction, chest tube, EVD, mechanical ventilation) 	<ul style="list-style-type: none"> <input type="checkbox"/> Yes = High Risk <input type="checkbox"/> No = Moderate Risk
<p>0-1 Risk Factor</p>	<p>Anticipated altered mobility > 48 hours?</p> <ul style="list-style-type: none"> <input type="checkbox"/> Bedrest or significant activity restriction <input type="checkbox"/> Any line or tube that restricts mobility (i.e. epidural, foley, NG to continuous suction, chest tube, EVD, mechanical ventilation) 	<ul style="list-style-type: none"> <input type="checkbox"/> Yes = Moderate Risk <input type="checkbox"/> No = Low Risk


If NO contraindications (see page 1), the following interventions (and orders) are indicated:

Pre-op Orders	
	Recommended Intervention/Order
<p><u>ALL</u> patients with surgery scheduled for > 60 minutes</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Intra-op SCD (sequential compression device) – apply in pre-op/holding area

Post-op or Admission Orders	
Risk Category	Recommended Interventions/Orders
<p>Low Risk</p>	<p>None – encourage early ambulation</p>
<p>Moderate Risk</p>	<ul style="list-style-type: none"> <input type="checkbox"/> SCD (sequential compression device), aim at 18 hours of use
<p>High Risk</p>	<ul style="list-style-type: none"> <input type="checkbox"/> SCD (sequential compression device), aim at 18 hours of use AND <input type="checkbox"/> Lovenox * (first dose 12 hours after surgery and hold 12h prior to surgical procedure) <ul style="list-style-type: none"> <input type="checkbox"/> < 50 kg = 0.5 mg/kg/dose subQ BID <input type="checkbox"/> 50-125 kg = 30 mg subQ BID <u>or</u> 40mg subQ daily <input type="checkbox"/> > 125 kg = 40 mg subQ BID <p>* If renal dysfunction, consider decreasing dose and checking LMWH level 4 hours after 2nd or 3rd dose (goal 0.1-0.3 unit/mL, see BEST statement for management of LMWH for more details) ** If considering other options, consult hematology.</p>

Algorithm: Risk Category Assessment and Prophylaxis for Venous Thromboembolism

Definitions, abbreviations and development credit for the algorithm – See next page



- Exclude patients:**
- 18 years and older (eligible for assessment and treatment by adult guideline)
 - with current VTE

*VTE Risk Factors
Blood stream infection
Central Venous Catheter (including non-tunneled, tunneled and PICCs)
History of venous thrombosis
Hyperosmolar state (serum osmolality >320 mOsm/kg)
Inflammatory diseases (e.g. IBD, SLE)
Medications: asparaginase, estrogen use (within past 2 months)
Obesity (BMI > 95 th percentile for age)
Oncologic diagnosis
Orthopedic procedures: hip or knee reconstruction
Nephrotic syndrome
Thrombophilia – known, or family history of clots
Trauma: >1 lower extremity long bone fracture, complex pelvic fractures, spinal cord injury
**Contraindications to Mechanical Prophylaxis
DVT, suspected or existing (can use GCS)
Extremity to be used has acute fracture
Extremity to be used has PIV access
Skin conditions affecting extremity (e.g. dermatitis, burn)
Unable to achieve correct fit due to patient size
***Contraindications to Anticoagulation
Absolute:
Bleeding disorder, known or tendency
Hemorrhage, evidence of or high risk of
Platelet count unable to be sustained > 50,000/mm ³
Relative:
Intracranial mass
Lumbar puncture or epidural catheter removal in prior 12 hours
Neurosurgical procedure
Pelvic fracture within past 48 hours
Uncontrolled hypertension

[Best Evidence Statement \(BEST\): Management of Low Molecular Weight Heparin Therapy \(LMWH\)](#)

If decision to initiate pharmacologic prophylaxis:

- in surgical patients seek surgical input regarding bleeding risk, prior to initiation
- see [BEST](#) for management of LMWH prophylaxis
- obtain hematology consult when considering alternative pharmacologic agents

Reassess risk at 48 to 72 hours of hospitalization

4.0 REFERENCES

4.1 Multidisciplinary VTE Prophylaxis BEST Team, Cincinnati Children's Hospital Medical Center: Best Evidence Statement Venous Thromboembolism (VTE) Prophylaxis in Children and Adolescents, <http://www.cincinnatichildrens.org/service/j/anderson-center/evidence-based-care/bests/>, BEST 181, pages 1-14, Date 2/18/14.

5.0 APPROVALS

All revisions of this guideline are approved by the Trauma Service Department. This guideline is reviewed every three years or sooner if deemed necessary. Policy authority for this document resides with the Trauma Service Department. This guideline is approved by the Trauma Services Manager and the Director of Trauma Services.

HISTORY	
Original Date	06/2004
Revision Date	05/15, 06/18
Review Date	